(1996). Luminescent β-galactosidase Genetic Reporter System II. *CLONTECHniques* **XI**(2), 16-17.

Beetham, J. K., Myung, K. S., McCoy, J. J., Wilson, M. E. & Donelson, J. E. (1997). Glycoprotein 46 mRNA abundance is post-transcriptionally regulated during development of *Leishmania chagasi* promastigotes to an infectious form. *J. Biol. Chem.* **272**(28), 17360-17366.

Besancon, F., Atfi, A., Gespach, C., Cayre, Y. E. & Bourgeade, M. F. (1998). Evidence for a role of NF-κB in the survival of hematopoietic cells mediated by interleukin 3 and the oncogenic TEL/platelet-derived growth factor receptor β fusion protein. *Proc. Natl. Acad. Sci. USA* **95**, 8081-8086.

Daub, M., Jockel, J., Quack, T., Weber, C. K., Schmitz, F., Rapp, U. R., Wittinghofer, A. & Block, C. (1998). The RafC1 cysteine-rich domain contains multiple distinct regulatory epitopes which control Ras-dependent Raf activation. *Mol. Cell. Biol.* **18**, 6698-6710.

Esser, K., Nelson, T., Lupa-Kimball, V. & Blough, E. (1999). The CACC box and myocyte enhancer factor-2 sites within the myosin light chain 2 slow promoter cooperate in regulating nerve-specific transcription in skeletal muscle. *J. Biol. Chem.* **274**, 12095-12102.

Lopez, R. G., Carron, C., Oury, C., Gardellin, P., Bernard, O., Ghysdael, J. (1999). TEL is a sequence-specific transcriptional repressor. *J. Biol. Chem.* **274:**30132-30138.

Martin, C. S., Olesen, C. E. M., Liu, B., Voyta, J. C., Shumway, J. L., Juo, R.-R. & Bronstein, I. (1997). Continuous sensitive detection of β-galactosidase with a novel chemiluminescent 1,2-dioxetane. In *Bioluminescence and Chemiluminescence: Molecular Reporting with Photons* (Hastings, J. W., Kricka, L. J. & Stanley, P. E., eds.), pp. 525-528. John Wiley, Chichester, England.

Olesen, C. E. M., Shumway, J. L., Martin, C. S. & Juo, R. R. (1996). Galacton-*Star*®: New chemiluminescent β-galactosidase substrate with "glow" kinetics. *Lumin. Topics* **5**(1), 1-7.

Olesnicky, N. S., Brown, A. J., Dowell, S. J. & Casselton, L. A. (1999). A constitutively active G-protein-coupled receptor causes mating self-compatibility in the mushroom *Coprinus*. *EMBO J.* **18**, 2756-2763.

Scharf, K. D., Heider, H., Hohfeld, I., Lyck, R., Schmidt, E. & Nover, L. (1998). The tomato Hsf system: HsfA2 needs interaction with HsfA1 for efficient nuclear import and may be localized in cytoplasmic heat stress granules. *Mol. Cell. Biol.* **18**(4), 2240-2251.

Weiland, P. R., Liu, B., Martin, C. S. & Olesen, C. E. M. (1997). Simplified detection of reporter gene expression using Galacto-*Star*™ β-galactosidase reporter gene assay system. *Lumin. Topics* **6**(1), 2.